

Problem Set 5

Problem 1 is required. All other problems are for your own practice.

1. (*Required problem*) Suppose that the random variables Y_1, \dots, Y_n satisfy

$$y_i = \beta x_i + e_i, \quad i = 1, \dots, n,$$

where x_1, \dots, x_n are fixed constants and e_1, \dots, e_n are i.i.d. normals with mean 0 and unknown variance σ^2 . Assume that the hypothesis of interest is $H_0 : \beta = 0$.

- Write the likelihood function (treating both β and σ^2 as unknown). Write down score and information matrix.
- Find the unrestricted maximum likelihood estimator. Write the Wald test for the null hypothesis.
- Solve the restricted maximization problem. Write the Lagrange Multiplier test.
- Write down the LR test.
- Introduce sample correlation between y_i and x_i :

$$\hat{r} = \frac{\sum_{i=1}^n y_i x_i}{\sqrt{\sum_{i=1}^n y_i^2} \sqrt{\sum_{i=1}^n x_i^2}}.$$

Re-write the Wald, LM and LR statistics as functions of \hat{r}^2 and n only.

- Notice, that for any $x < 1$ the following inequality holds: $x < -\ln(1-x) < \frac{x}{1-x}$. This implies some ordering of the statistics discussed above. What is it? Apparently, it holds in general for linear hypothesis in OLS models.
2. Assume that n_1 people are given treatment 1 and n_2 people are given treatment 2. Let X_1 be the number of people on treatment 1 who respond favorably to

the treatment and let X_2 be the number of people on treatment 2 who respond favorably. Assume that $X_1 \sim \text{Binomial}(n_1, p_1)$, $X_2 \sim \text{Binomial}(n_2, p_2)$ and $n_1 = \gamma n$, $n_2 = (1 - \gamma)n$. Let $\psi = p_1 - p_2$.

- (a) Assume that the unknown parameters are (p_1, p_2) , write the likelihood. Find the MLE estimator of ψ . Would your answer change if you write likelihood in terms of parameters (ψ, p_2) ?
 - (b) Find the Fisher information matrix $I(p_1, p_2)$.
 - (c) Use the multiparameter delta-method to find the asymptotic variance of $\hat{\psi}$ assuming that $n \rightarrow \infty$.
 - (d) Construct a Wald confidence set for ψ . Is it asymptotic? Why or why not?
 - (e) Test the null hypothesis $\psi = 0$ using the asymptotic LR test. Describe all the details.
3. Suppose that X_1, \dots, X_n is a random sample from $N(\mu, \sigma^2)$ with known σ^2 . Find a minimum value of n to guarantee that a 0.95 confidence interval for μ will have length no more than $\frac{\sigma}{4}$.
4. Assume that X_1, \dots, X_n are iid Poisson (λ)
- (a) Construct a Wald type confidence set for λ .
 - (b) Construct a confidence set for λ by inverting Lagrange multiplier (score) test.

MIT OpenCourseWare
<https://ocw.mit.edu>

14.381 Statistical Method in Economics
Fall 2018

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>