



September 15, 2016

**6.453 Quantum Optical Communication**  
**Lecture 3**

Jeffrey H. Shapiro

Optical and Quantum Communications Group

RESEARCH LABORATORY OF ELECTRONICS  
Massachusetts Institute of Technology

[www.rle.mit.edu/qoptics](http://www.rle.mit.edu/qoptics)

### **6.453 Quantum Optical Communication — Lecture 3**

- Announcements
  - Turn in problem set 1
  - Pick up problem set 1 solution, problem set 2, lecture notes, slides
- Fundamentals of Dirac-Notation Quantum Mechanics
  - Definitions and axioms — reprise
  - Quantum measurements — statistics
  - Schrödinger picture versus Heisenberg picture
  - Heisenberg uncertainty principle

## Quantum Systems and Quantum States

- **Definition 1:**

A quantum-mechanical system  $\mathcal{S}$  is a physical system governed by the laws of quantum mechanics.

- **Definition 2:**

The state of a quantum mechanical system at a particular time  $t$  is the sum total of all information that can be known about the system at time  $t$ . It is a ket vector  $|\psi(t)\rangle$  in an appropriate Hilbert space  $\mathcal{H}_{\mathcal{S}}$  of possible states. Finite energy states have unit length ket vectors, i.e.,  $\langle\psi(t)|\psi(t)\rangle = 1$ .

## Time Evolution via the Schrödinger Equation

- **Axiom 1:**

For  $t \geq 0$ , an isolated system with initial state  $|\psi(0)\rangle$  will reach state

$$|\psi(t)\rangle = \hat{U}(t, 0)|\psi(0)\rangle$$

where  $\hat{U}(t, 0)$  is the unitary time-evolution operator for the system  $\mathcal{S}$ .  $\hat{U}(t, 0)$  is obtained by solving

$$j\hbar \frac{d\hat{U}(t, 0)}{dt} = \hat{H}\hat{U}(t, 0), \quad \text{for } t \geq 0, \text{ with } \hat{U}(0, 0) = \hat{I}$$

where  $\hat{H}$  is the Hamiltonian (energy) operator for  $\mathcal{S}$ . Equivalently, we have the Schrödinger equation

$$j\hbar \frac{d|\psi(t)\rangle}{dt} = \hat{H}|\psi(t)\rangle, \quad \text{for } t \geq 0, \text{ with } |\psi(0)\rangle \text{ initial condition}$$

## Quantum Measurements: Observables

- *Axiom 2:*

An observable is a measurable dynamical variable of the quantum system  $\mathcal{S}$ . It is represented by an Hermitian operator which has a complete set of eigenkets.

- *Axiom 3:*

For a quantum system  $\mathcal{S}$  that is in state  $|\psi(t)\rangle$  at time  $t$ , measurement of the observable

$$\hat{O} \equiv \sum_n o_n |o_n\rangle\langle o_n|$$

yields an outcome that is one of the eigenvalues,  $\{o_n\}$ , with

$$\Pr(\text{outcome} = o_n) = |\langle o_n | \psi(t) \rangle|^2$$

## Quantum Measurements: Observables

- *Projection postulate:*

Immediately after a measurement of an observable  $\hat{O}$ , with distinct eigenvalues, yields outcome  $o_n$  the state of the system becomes  $|o_n\rangle$ .

- *Axiom 3a:*

For a quantum system  $\mathcal{S}$  that is in state  $|\psi(t)\rangle$  at time  $t$ , measurement of the observable

$$\hat{O} = \int_{-\infty}^{\infty} do o |o\rangle\langle o|$$

yields an outcome that is one of the eigenvalues,  $o$ , with

$$p(o) = |\langle o | \psi(t) \rangle|^2$$

## Quantum Measurements: Statistics

- Average Value of an Observable Measurement
  - Discrete eigenvalue spectrum

$$\langle \hat{O} \rangle \equiv \sum_n o_n \Pr(\text{outcome} = o_n) = \sum_n o_n |\langle o_n | \psi(t) \rangle|^2 = \langle \psi(t) | \hat{O} | \psi(t) \rangle$$

- Continuous eigenvalue spectrum

$$\langle \hat{O} \rangle \equiv \int_{-\infty}^{\infty} do op(o) = \int_{-\infty}^{\infty} do |\langle o | \psi(t) \rangle|^2 = \langle \psi(t) | \hat{O} | \psi(t) \rangle$$

- Variance of an Observable Measurement

$$\langle \Delta \hat{O}^2 \rangle = \langle \psi(t) | \Delta \hat{O}^2 | \psi(t) \rangle, \quad \text{for } \Delta \hat{O} \equiv \hat{O} - \langle \hat{O} \rangle$$

## Schrödinger versus Heisenberg Pictures

- Schrödinger Picture
  - Observables are time-independent operators
  - Between measurements, states evolve according to the Schrödinger equation

$$\{ |\psi(t)\rangle_S, \hat{O}_S, \hat{H}_S : t \geq 0 \}$$

- Heisenberg Picture
  - Between measurements, states are constant
  - Observables evolve in time according to appropriate equations of motion

$$\{ |\psi\rangle_H, \hat{O}_H(t), \hat{H}_H(t) : t \geq 0 \}$$

## Converting Between Pictures

- Statistics of an Observable Measurement
  - Schrödinger picture

$$\Pr(\text{outcome} = o_n) = |{}_S\langle o_n | \psi(t) \rangle_S|^2$$

- Heisenberg picture

$$\Pr(\text{outcome} = o_n) = |{}_H\langle o_n(t) | \psi \rangle_H|^2$$

- Invariance of Statistics to Choice of Picture

$$|o_n(t)\rangle_H = \hat{U}^\dagger(t, 0)|o_n\rangle_S$$

## Heisenberg Equations of Motion

- Transforming an Observable between Pictures

$$\hat{O}_H(t) = \hat{U}^\dagger(t, 0)\hat{O}_S\hat{U}(t, 0)$$

- Equation of Motion for  $\hat{O}_H(t)$

$$j\hbar \frac{d\hat{O}_H(t)}{dt} = [\hat{O}_H(t), \hat{H}], \quad \text{for } t \geq 0, \text{ with } \hat{O}_H(0) = \hat{O}_S$$

- Commutator Brackets

$$[\hat{O}_H(t), \hat{H}] \equiv \hat{O}_H(t)\hat{H} - \hat{H}\hat{O}_H(t)$$

## Heisenberg Uncertainty Principle

- $\hat{A}(t)$  and  $\hat{B}(t)$  Noncommuting Observables

$$[\hat{A}(t), \hat{B}(t)] = j\hat{C}(t), \quad \hat{C}(t) \text{ is Hermitian}$$

- Lower Limit on Product of Individual Measurement Variances

$$\langle \Delta \hat{A}(t)^2 \rangle \langle \Delta \hat{B}(t)^2 \rangle \geq \frac{1}{4} |\langle \hat{C}(t) \rangle|^2$$

## Coming Attractions: Lectures 4 and 5

- Lecture 4:  
Quantum Harmonic Oscillator
  - Quantization of a classical  $LC$  circuit
  - Annihilation and creation operators
  - Energy eigenstates — number-state kets
- Lecture 5:  
Quantum Harmonic Oscillator
  - Quadrature measurements
  - Coherent states

MIT OpenCourseWare  
<https://ocw.mit.edu>

6.453 Quantum Optical Communication  
Fall 2016

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.