

**THIRD PRACTICE MIDTERM
MATH 18.022, MIT, AUTUMN 10**

You have 50 minutes. This test is closed book, closed notes, no calculators.

Name: MODEL ANSWERS

Signature: _____

Recitation Time: _____

There are 5 problems, and the total number of points is 100. Show all your work. *Please make your work as clear and easy to follow as possible.*

Problem	Points	Score
1	20	
2	20	
3	20	
4	20	
5	20	
Total	100	

1. (20pts) For what values of λ , μ and ν does the function $f: \mathbb{R}^3 \rightarrow \mathbb{R}$,

$$f(x, y, z) = \lambda x^2 + \mu xy + y^2 + \nu z^2,$$

have a non-degenerate local minimum at $(0, 0, 0)$?

$$Df = (2\lambda x + \mu y, 2y + \mu x, 2\nu z)$$

$$Hf = \begin{pmatrix} 2\lambda & \mu & 0 \\ \mu & 2 & 0 \\ 0 & 0 & 2\nu \end{pmatrix}$$

$$d_1 = 2\lambda, \quad d_2 = 4\lambda - \mu^2, \quad d_3 = 2\nu \cdot d_2.$$

Minimum: $d_1 > 0, d_2 > 0, d_3 > 0.$

So $\lambda \geq 0, \mu^2 < 4\lambda, \nu \geq 0$

2. (20pts) Let $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ be the function $f(x, y, z) = 2x + y - z$

(i) Show that f has a global minimum on the ellipsoid $x^2 + 2y^2 + 3z^2 = 6$.

$$K = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + 2y^2 + 3z^2 = 6\}$$

is closed + bounded. So K is compact.

f is cts, K is compact $\Rightarrow f$ has a global minimum.

(ii) Find this minimum.

Consider $h: \mathbb{R}^4 \rightarrow \mathbb{R}$ given by

$$h(x, y, z, \lambda) = 2x + y - z - \lambda(x^2 + 2y^2 + 3z^2 - 6)$$

Critical pts of h :

$$x = \frac{1}{\lambda} \quad x = 4y = -6z$$

$$4y = \frac{1}{\lambda} \quad y = -\frac{3z}{2}, \quad x = -6z.$$

$$-6z = \frac{1}{\lambda}$$

$$x^2 + 2y^2 + 3z^2 = 6 \quad z^2 \left(3 + \frac{9}{2} + 36 \right) = 6$$

$$z^2 (2 + 3 + \frac{24}{2}) = 4$$

$$z = \frac{\pm \sqrt{29}}{\sqrt{29}}$$

$$f(P) = \left(24 + 3 + 2 \right) \frac{1}{\sqrt{29}}$$

min value: + square root


$$P = \left(\frac{-12}{\sqrt{29}}, \frac{-3}{\sqrt{29}}, \frac{2}{\sqrt{29}} \right)$$

$$= -\frac{2}{\sqrt{29}}$$

3. (20pts)

(i) Draw a picture of the region of integration of

$$\int_0^1 \int_{1+x}^{\sqrt{9-x^2}} dy dx.$$



(ii) Change the order of integration of the integral.

$$\int_1^2 \int_0^{y-1} dx dy + \int_2^{\sqrt{8}} \int_0^1 dx dy + \int_{\sqrt{8}}^3 \int_0^{\sqrt{9-y^2}} dx dy$$

4. (20pts) Let W be the region inside the two cylinders $x^2 + y^2 = 1$ and $y^2 + z^2 = 1$.

Set up an integral to calculate the volume of W and calculate this integral.

View as a region of type 2.

$$\begin{aligned}\text{vol}(W) &= \iiint dx dy dz \\ &= \int_{-1}^1 \left(\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \left(\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} dx \right) dz \right) dy \\ &= 2 \int_{-1}^1 \left(\int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \sqrt{1-y^2} dz \right) dy \\ &= 4 \int_{-1}^1 (1-y^2) dy \\ &= 4 \left[y - \frac{y^3}{3} \right]_{-1}^1 \\ &= 8 \left(1 - \frac{1}{3} \right) = \frac{16}{3}.\end{aligned}$$

5. (20pts) Let D be the region in the first quadrant bounded by the curves $y^2 = x$, $y^2 = 2x$, $xy = 1$ and $xy = 4$.

(i) Find $du dv$ in terms of $dx dy$, where $u = \frac{y^2}{x}$ and $v = xy$.

$$\frac{\partial(u,v)}{\partial(x,y)} = \begin{vmatrix} -\frac{y^2}{x^2} & \frac{2y}{x} \\ y & x \end{vmatrix} = -\frac{y^2}{x} - \frac{2y^2}{x} = -\frac{3y^2}{x}$$

$$\frac{\partial(x,y)}{\partial(u,v)} = \frac{-x}{3y^2} \quad dx dy = (3u)^{-1} du dv$$

(ii) Set up an integral to calculate the area of the region D and calculate this integral.

$$\begin{aligned} \iint_D dx dy &= \int_1^4 \left(\int_1^2 \frac{du}{3u} \right) dv \\ &= \frac{1}{3} \int_1^4 [\ln u]_1^2 dv \\ &= \frac{\ln 2}{3} \int_1^4 dv \\ &= \ln 2. \end{aligned}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

18.022 Calculus of Several Variables
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.