

## Recitation 13, March 18, 2010

### Fourier Series: Introduction

1. What is the general solution to  $\ddot{x} + \omega_n^2 x = 0$ ? [Quick!]
2. Discuss why (as long as  $\omega \neq \pm\omega_n$ )

$$\ddot{x} + \omega_n^2 x = a \cos(\omega t) \quad \text{has solution} \quad x_p = a \frac{\cos(\omega t)}{\omega_n^2 - \omega^2}$$

$$\ddot{x} + \omega_n^2 x = b \sin(\omega t) \quad \text{has solution} \quad x_p = b \frac{\sin(\omega t)}{\omega_n^2 - \omega^2}$$

3. What about  $\ddot{x} + \omega_n^2 x = \cos(\omega_n t)$ ? What is a particular solution? What is the general solution? Are there any solutions  $x(t)$  such that  $|x(t)| < 10^6$  for all  $t$ ? Are there any periodic solutions?

A function is *periodic* if there is a number  $P > 0$  such that  $f(t+P) = f(t)$  for all  $t$ . Such a number  $P$  is then a “period” of  $f(t)$ . If  $f(t)$  is a periodic function which is continuous and not constant, then there is a smallest period, often called *the* period.

4. On the same set of axes, sketch graphs of  $\sin(t)$ ,  $\sin(2t)$ . Then sketch the graph of  $f(t) = \sin(t) + \sin(2t)$ . Some pointers:  $f(t)$  is easy to evaluate when one of the terms is zero. What is the derivative at points where both terms are zero? This information should be enough to let you make a rough sketch. What are the periods of these three functions?

5. For what values of  $\omega_n$  is there a periodic solution to the equation

$$\ddot{x} + \omega_n^2 x = b_1 \sin(t) + b_2 \sin(2t)$$

(where  $b_1$  and  $b_2$  are nonzero)? Name one if it exists.

6. (very tricky) For what values of  $\omega$  is  $\sin(t) + \sin(\omega t)$  periodic? And the periods?

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.03 Differential Equations  
Spring 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.