


# Bayesian Updating: Discrete Priors: 18.05 Spring 2014



STATISTICALLY SPEAKING, IF YOU PICK UP A SEASHELL AND DON'T HOLD IT TO YOUR EAR, YOU CAN PROBABLY HEAR THE OCEAN.

<http://xkcd.com/1236/>

## Learning from experience

Which treatment would you choose?

1. Treatment 1: cured 100% of patients in a trial.
2. Treatment 2: cured 95% of patients in a trial.
3. Treatment 3: cured 90% of patients in a trial.

Which treatment would you choose?

1. Treatment 1: cured 3 out of 3 patients in a trial.
2. Treatment 2: cured 19 out of 20 patients treated in a trial.
3. Standard treatment: cured 90000 out of 100000 patients in clinical practice.

## Which die is it?

- I have a bag containing dice of two types: 4-sided and 10-sided.
  - Suppose I pick a die at random and roll it.
  - Based on what I rolled which type would you guess I picked?
- 
- Suppose you find out that the bag contained one 4-sided die and one 10-sided die. Does this change your guess?
- 
- Suppose you find out that the bag contained one 4-sided die and 100 10-sided dice. Does this change your guess?

## Board Question: learning from data

- A certain disease has a prevalence of 0.005.
- A screening test has 2% false positives and 1% false negatives.

Suppose a patient is screened and has a positive test.

- 1 Represent this information with a tree and use Bayes' theorem to compute the probabilities the patient does and doesn't have the disease.
- 2 Identify the data, hypotheses, likelihoods, prior probabilities and posterior probabilities.
- 3 Make a full **likelihood table** containing all hypotheses and possible test data.
- 4 Redo the computation using a **Bayesian update table**. Match the terms in your table to the terms in your previous calculation.

## Board Question: Dice

Five dice: 4-sided, 6-sided, 8-sided, 12-sided, 20-sided.

Suppose I picked one at random and, without showing it to you, rolled it and reported a 13.

1. Make the full likelihood table (be smart about identical columns).
2. Make a Bayesian update table and compute the posterior probabilities that the chosen die is each of the five dice.
3. Same question if I rolled a 5.
4. Same question if I rolled a 9.

(Keep the tables for 5 and 9 handy! Do not erase!)



## Tabular solution

$\mathcal{D}$  = 'rolled a 13'

hypothesis	prior	likelihood	Bayes	
			numerator	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D} \mathcal{H})$	$P(\mathcal{D} \mathcal{H})P(\mathcal{H})$	$P(\mathcal{H} \mathcal{D})$
$\mathcal{H}_4$	1/5	0	0	0
$\mathcal{H}_6$	1/5	0	0	0
$\mathcal{H}_8$	1/5	0	0	0
$\mathcal{H}_{12}$	1/5	0	0	0
$\mathcal{H}_{20}$	1/5	1/20	1/100	1
total	1		1/100	1

## Tabular solution

$\mathcal{D}$  = 'rolled a 5'

hypothesis	prior	likelihood	Bayes	
			numerator	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D} \mathcal{H})$	$P(\mathcal{D} \mathcal{H})P(\mathcal{H})$	$P(\mathcal{H} \mathcal{D})$
$\mathcal{H}_4$	1/5	0	0	0
$\mathcal{H}_6$	1/5	1/6	1/30	0.392
$\mathcal{H}_8$	1/5	1/8	1/40	0.294
$\mathcal{H}_{12}$	1/5	1/12	1/60	0.196
$\mathcal{H}_{20}$	1/5	1/20	1/100	0.118
total	1		0.085	1

## Tabular solution

$\mathcal{D}$  = 'rolled a 9'

hypothesis	prior	likelihood	Bayes	
			numerator	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D} \mathcal{H})$	$P(\mathcal{D} \mathcal{H})P(\mathcal{H})$	$P(\mathcal{H} \mathcal{D})$
$\mathcal{H}_4$	1/5	0	0	0
$\mathcal{H}_6$	1/5	0	0	0
$\mathcal{H}_8$	1/5	0	0	0
$\mathcal{H}_{12}$	1/5	1/12	1/60	0.625
$\mathcal{H}_{20}$	1/5	1/20	1/100	0.375
total	1		.0267	1



# Iterated Updates

Suppose I rolled a 5 and then a 9.

Update in two steps:

First for the 5

Then update the update for the 9.

## Tabular solution

$\mathcal{D}_1 =$  'rolled a 5'

$\mathcal{D}_2 =$  'rolled a 9'

Bayes numerator<sub>1</sub> = likelihood<sub>1</sub> × prior.

Bayes numerator<sub>2</sub> = likelihood<sub>2</sub> × Bayes numerator<sub>1</sub>

hyp.	prior	likel. 1	Bayes num. 1	likel. 2	Bayes num. 2	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D}_1 \mathcal{H})$	***	$P(\mathcal{D}_2 \mathcal{H})$	***	$P(\mathcal{H} \mathcal{D}_1, \mathcal{D}_2)$
$\mathcal{H}_4$	1/5	0	0	0	0	0
$\mathcal{H}_6$	1/5	1/6	1/30	0	0	0
$\mathcal{H}_8$	1/5	1/8	1/40	0	0	0
$\mathcal{H}_{12}$	1/5	1/12	1/60	1/12	1/720	0.735
$\mathcal{H}_{20}$	1/5	1/20	1/100	1/20	1/2000	0.265
total	1				0.0019	1

## Board Question

Suppose I rolled a 9 and then a 5.

1. Do the Bayesian update in two steps:  
First update for the 9.  
Then update the update for the 5.
2. Do the Bayesian update in one step  
The data is  $\mathcal{D} = \text{'9 followed by 5'}$

## Tabular solution: two steps

$\mathcal{D}_1 =$  'rolled a 9'

$\mathcal{D}_2 =$  'rolled a 5'

Bayes numerator<sub>1</sub> = likelihood<sub>1</sub> × prior.

Bayes numerator<sub>2</sub> = likelihood<sub>2</sub> × Bayes numerator<sub>1</sub>

hyp.	prior	likel. 1	Bayes num. 1	likel. 2	Bayes num. 2	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D}_1 \mathcal{H})$	***	$P(\mathcal{D}_2 \mathcal{H})$	***	$P(\mathcal{H} \mathcal{D}_1, \mathcal{D}_2)$
$\mathcal{H}_4$	1/5	0	0	0	0	0
$\mathcal{H}_6$	1/5	0	0	1/6	0	0
$\mathcal{H}_8$	1/5	0	0	1/8	0	0
$\mathcal{H}_{12}$	1/5	1/12	1/60	1/12	1/720	0.735
$\mathcal{H}_{20}$	1/5	1/20	1/100	1/20	1/2000	0.265
total	1				0.0019	1

## Tabular solution: one step

$\mathcal{D}$  = 'rolled a 9 then a 5'

hypothesis	prior	likelihood	Bayes	
			numerator	posterior
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D} \mathcal{H})$	$P(\mathcal{D} \mathcal{H})P(\mathcal{H})$	$P(\mathcal{H} \mathcal{D})$
$\mathcal{H}_4$	1/5	0	0	0
$\mathcal{H}_6$	1/5	0	0	0
$\mathcal{H}_8$	1/5	0	0	0
$\mathcal{H}_{12}$	1/5	1/144	1/720	0.735
$\mathcal{H}_{20}$	1/5	1/400	1/2000	0.265
total	1		0.0019	1

## Board Question: probabilistic prediction

Along with finding posterior probabilities of hypotheses. We might want to make **posterior predictions** about the next roll.

With the same setup as before let:

$\mathcal{D}_1$  = result of first roll

$\mathcal{D}_2$  = result of second roll

(a) Find  $P(\mathcal{D}_1 = 5)$ .

(b) Find  $P(\mathcal{D}_2 = 4 | \mathcal{D}_1 = 5)$ .

## Solution

$\mathcal{D}_1 =$  'rolled a 5'

$\mathcal{D}_2 =$  'rolled a 4'

hyp.	prior	likel. 1	Bayes			
			num. 1	post. 1	likel. 2	post. 1 $\times$ likel. 2
$\mathcal{H}$	$P(\mathcal{H})$	$P(\mathcal{D}_1 \mathcal{H})$	***	$P(\mathcal{H} \mathcal{D}_1)$	$P(\mathcal{D}_2 \mathcal{H}, \mathcal{D}_1)$	$P(\mathcal{D}_2 \mathcal{H}, \mathcal{D}_1)P(\mathcal{H} \mathcal{D}_1)$
$\mathcal{H}_4$	1/5	0	0	0	*	0
$\mathcal{H}_6$	1/5	1/6	1/30	0.392	1/6	0.392 $\cdot$ 1/6
$\mathcal{H}_8$	1/5	1/8	1/40	0.294	1/8	0.294 $\cdot$ 1/40
$\mathcal{H}_{12}$	1/5	1/12	1/60	0.196	1/12	0.196 $\cdot$ 1/12
$\mathcal{H}_{20}$	1/5	1/20	1/100	0.118	1/20	0.118 $\cdot$ 1/20
total	1		0.085	1		0.124

The law of total probability tells us  $P(\mathcal{D}_1)$  is the sum of the Bayes numerator 1 column in the table:  $P(\mathcal{D}_1) = 0.085$ .

The law of total probability tells us  $P(\mathcal{D}_2|\mathcal{D}_1)$  is the sum of the last column in the table:  $P(\mathcal{D}_2|\mathcal{D}_1) = 0.124$

MIT OpenCourseWare  
<https://ocw.mit.edu>

## 18.05 Introduction to Probability and Statistics

Spring 2014

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.